

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/291184680>

The fecal virome of South and Central American children with diarrhea includes small circular DNA viral genomes of unknown origin

ARTICLE *in* ARCHIVES OF VIROLOGY · JANUARY 2016

Impact Factor: 2.39 · DOI: 10.1007/s00705-016-2756-4

READS

70

8 AUTHORS, INCLUDING:



[Antonio Charlys Da-Costa](#)

University of São Paulo

14 PUBLICATIONS 40 CITATIONS

[SEE PROFILE](#)



[Filemon Bucardo](#)

National Autonomous University of Nicarag...

31 PUBLICATIONS 418 CITATIONS

[SEE PROFILE](#)



[Johan Nordgren](#)

Linköping University

41 PUBLICATIONS 541 CITATIONS

[SEE PROFILE](#)



[Eric Delwart](#)

University of California, San Francisco

240 PUBLICATIONS 11,286 CITATIONS

[SEE PROFILE](#)

The fecal virome of South and Central American children with diarrhea includes small circular DNA viral genomes of unknown origin

Tung Gia Phan^{1,2} · Antonio Charlys da Costa^{1,3} · Juana del Valle Mendoza^{4,5} · Filemon Bucardo-Rivera⁶ · Johan Nordgren⁷ · Miguel O’Ryan⁸ · Xutao Deng¹ · Eric Delwart^{1,2}

Received: 13 October 2015 / Accepted: 8 January 2016
© Springer-Verlag Wien 2016

Abstract Viral metagenomics of feces collected from 58 Peruvian children with unexplained diarrhea revealed several small circular ssDNA genomes. Two genomes related to sequences previously reported in feces from chimpanzees and other mammals and recently named smacoviruses were characterized and then detected by PCR in 1.7 % (1/58) and 19 % (11/58) of diarrheal samples, respectively. Another three genomes from a distinct small circular ssDNA viral group provisionally called pecoviruses encoded Cap and Rep proteins with <35 % identity to those in related genomes reported in human, seal, porcine and dromedary feces. Pecovirus DNA was detected in 15.5 % (9/58), 5.9 % (3/51) and 3 % (3/100) of fecal samples from unexplained diarrhea

in Peru, Nicaragua and Chile, respectively. Feces containing these ssDNA genomes also contained known human enteric viral pathogens. The cellular origins of these circular ssDNA viruses, whether human cells, ingested plants, animals or fungal foods, or residents of the gut microbiome, are currently unknown.

Introduction

The known diversity of small circular Rep-encoding ssDNA (CRESS-DNA) genomes has rapidly increased due largely to the analysis of environmental and animal samples using deep sequencing following rolling-circle amplification [6, 11–13, 20, 32]. Among the many distinct taxa of CRESS-DNA viruses, a few are known to replicate in eukaryotic hosts ranging from plants and fungi to birds and mammals [7]. The cellular hosts for many viruses with a CRESS-DNA genome remain unknown. Members of the family *Circoviridae* include circoviruses known to infect birds and mammals, at times with pathogenic consequences. Genomes closely related to those of circoviruses, named “cycloviruses”, have been detected in mammalian tissues and feces as well as in the bodies of different insects [7, 9, 19, 25, 31]. For cycloviruses, the cellular hosts remain unknown but may include mammals and/or insects cells, or conceivably, originate from consumed animals or plants. Another distinct clade of viruses with CRESS-DNA genomes was also described in the feces of chimpanzees and other mammals, including pigs, rats [1, 3–5, 10, 18, 21, 22, 25, 26, 30], and recently humans [14], whose cellular hosts are also unknown. Yet another clade of CRESS-DNA genomes whose cellular hosts are not known was reported in the feces of pigs, seals, dromedaries and humans [4, 24, 27, 30]. Here we describe five CRESS-DNA genomes in human fecal

✉ Eric Delwart
delwarte@medicine.ucsf.edu

¹ Blood Systems Research Institute, San Francisco, CA 94118, USA

² Department of Laboratory Medicine, University of California at San Francisco, San Francisco, CA 94118, USA

³ Institute of Tropical Medicine, School of Medicine, University of São Paulo, São Paulo, Brazil

⁴ Faculty of Health Sciences, School of Medicine, Universidad Peruana de Ciencias Aplicadas-UPC, Lima, Peru

⁵ Instituto de Investigación Nutricional, Lima, Peru

⁶ Department of Microbiology, University of León, León, Nicaragua

⁷ Division of Molecular Virology, Department of Clinical and Experimental Medicine, Linköping University, Linköping, Sweden

⁸ Microbiology and Mycology Program, Faculty of Medicine, University of Chile, Santiago, Chile

samples that cluster within two of these phylogenetic clades of unknown cellular origin. We characterize these genomes and describe their prevalence as measured by PCR in human diarrhea fecal samples.

Materials and methods

Clinical samples and viral metagenomics

Fecal samples from Peruvian children with diarrhea of unknown etiology were used that pre-tested negative for rotavirus, adenovirus, norovirus, campylobacter, shigella, salmonella and *Escherichia coli* [8]. A total of 58 fecal samples were analyzed in pools of three to five samples. Fecal suspensions were first vortexed and clarified by centrifugation at 15,000×g for 10 min. The supernatant was filtered through a 0.45-µm filter (Millipore) to remove bacterium-sized particles. The filtrate was treated with a mixture of enzymes to digest unprotected nucleic acids [16], and viral nucleic acids were then extracted using a Maxwell 16 automated extractor (Promega). Random RT-PCR using primers with degenerate 3' ends was then used to amplify RNA and DNA molecules, and a library was constructed for Illumina sequencing (MiSeq 2x250 bases) using a Nextera™ XT Sample Preparation Kit. Translated sequence reads showing similarity to viral sequences with an E score <10⁻⁵ were identified using BLASTx. Viral sequences were identified based on similarity to annotated viral genome sequences available in the GenBank non-redundant database, and results were mapped to the NCBI Taxonomy.

Determination of genome sequences of novel circular viruses

Complete circular DNA viral genomes were amplified using inverse PCR (iPCR) with specific primers designed from short sequence fragments. iPCR amplicons were then sequenced directly by primer walking. Putative ORFs in the genome were predicted using NCBI ORF Finder. Sequence analysis was performed using ClustalX with default settings [23]. A sequence identity matrix was made using BioEdit. To identify stem-loop structures, nucleotide sequences were analyzed with Mfold [33]. Phylogenetic trees with 100 bootstrap resamplings of the alignment data sets were generated using the neighbor-joining method and visualized using the program MEGA version 6 [28]. Bootstrap values (based on 100 replicates) for each node are shown if >70 %.

PCR assays for novel circular viruses

Three sets of primers for nested PCR based on the Rep coding regions were designed to screen nucleic acids from fecal supernatants.

For CRESS-DNA genome PE31, primers PESmaCV-F1 (5'-GGA TGT TAT TCG GTT GTG CGG A-3') and PESmaCV-R1 (5'-CAC CGC AGA ATG ATC CGA CAA-3') were used for the first round of PCR, and primers PESmaCV-F2 (5'-CAG TCC GTC TTT GAT GCT TTC-3') and PESmaCV-R2 (5'-GGA GTA TGA GGC GAA AGA AGG-3') were used for the second round of PCR, resulting in an expected amplicon of 255 bp. The PCR conditions were 95 °C for 5 min, 35 cycles of 95 °C for 30 s, 52 °C or 50 °C (for the first and second round, respectively) for 30 s and 72 °C for 1 min, followed by a final extension at 72 °C for 10 min.

For CRESS-DNA genome PE16, primers PESmaCV-F3 (5'-CTT TGG TTG TGT GTT CGT TAG GAC T-3') and PESmaCV-R3 (5'-AGC AGC AAC ACA GGA TTC TTC-3') were used for the first round of PCR, and primers PESmaCV-F4 (5'-TGG TAT CTC GTG TCG TAA ACT-3') and PESmaCV-R4 (5'-GAA AAA GCC GAA AGA GGA GTG-3') were used for the second round of PCR, resulting in an expected amplicon of 425 bp. The PCR conditions were 95 °C for 5 min, 35 cycles of 95 °C for 30 s, 50 °C or 49 °C (for the first and second round, respectively) for 30 s and 72 °C for 1 min, followed by a final extension at 72 °C for 10 min.

For CRESS-DNA genome PeCV-PE, primers PEPeCV-F1 (5'-TCC TTC ATC GTG GTA GCA ATG GTA G-3') and PEPeCV-R1 (5'-CTG GTG TGT AAC GAG TTG GAC TTG C-3') were used for the first round of PCR, and primers PEPeCV-F2 (5'-CTC ACG CTG TTT AGC CCT TTT G-3') and PEPeCV-R2 (5'-TGA TAG CGT GGA ACG TCT TGT AGT-3') were used for the second round of PCR, resulting in an expected amplicon of 430 bp. The PCR conditions were 95 °C for 5 min, 35 cycles of 95 °C for 30 s, 53 °C or 51 °C (for the first and second round, respectively) for 30 s and 72 °C for 1 min, followed by a final extension at 72 °C for 10 min.

Results

Viral metagenomics overview

Viral-particle-associated nucleic acids were enriched and randomly amplified from 58 fecal samples from Peruvian children with diarrhea of unknown etiology. Thirty-six

million sequence reads were generated in one Illumina MiSeq run. Using a BLASTx E-score cutoff of 10^{-5} , sequence reads related to mammalian viruses consisted of norovirus (124,168 reads), adenovirus (86,714 reads), group A rotavirus (35,869 reads), anelloviruses (21,995 reads), enterovirus (4,670 reads), CRESS-DNA-like viruses (3,043 reads), salivirus (461 reads), group C rotavirus (198 reads), bocavirus (93 reads), picobirnavirus (74 reads), sapovirus (73 reads) and Aichi virus (2 reads).

New clade distantly related to members of the *Circoviridae*




One fecal sample pool of five samples contained 1,753 sequence reads sharing low levels of genetic similarity to the Rep and Cap proteins of circular ssDNA viruses found in feces from a human (GenBank KJ206566), a pig (GenBank KJ414134) and a seal (GenBank KF246569). The individual fecal sample within the pool containing these sequences was identified using PCR. Using inverse PCR and Sanger sequencing, we then determined the complete genome sequence of the virus. To facilitate description and discussion of that clade, we provisionally refer to that group of viral genomes as “pecoviruses” (Peruvian stool-associated circo-like viruses). The circular human-feces-derived pecovirus genome (PeCV-PE) contained 2,937 bases, with a GC content of 49 %, and two major ORFs encoding replicase and capsid proteins (GenBank KT600065). Similar to many CRESS-DNA genomes, the Cap and Rep genes were in opposite orientation [19]. The Rep and Cap proteins were 320 aa and 388 aa in length, respectively. The Rep showed 35 % aa sequence identity to Rep of porcine- and human-feces-derived CRESS-DNA genomes. A Rep-based phylogenetic analysis showed that PeCV-PE clustered together with previously reported CRESS-DNA genomes from human, seal, pig, and dromedary feces. While the genomes from animal feces all shared the highly conserved canonical nonamer (NAN-TATTAC) atop a stem-loop, the conserved nonamer was not found in the human-feces-derived PeCV genomes, which contained a stem-loop with the nonamer TTTTAT-GAG. Direct observation of the PeCV Rep sequence alignment revealed two rolling-circle replication motifs II (xHxH) and III (YxxK) [19]. The seal-feces-derived PeCV contained a rolling-circle replication motif I (LTVKN) that was not found in other PeCV genomes [24]. Similarly, the C-terminal region of the PeCV Rep protein possessed the ATP-dependent helicase Walker A (GxxGxGKS) and Walker B (IWFDEFNG) motifs [19, 24]. The Walker C

motif (UxxN) was not found in any of the PeCV genomes [19, 24]. Pairwise comparison of all available PeCV sequences from the feces of humans, seals, pigs, and dromedaries demonstrated a high level of genetic diversity, with Rep identities of only 29–35 %, suggesting the existence of multiple species in that clade.

Nested PCR was then used to specifically test for human-associated PeCV DNA in all 58 fecal samples from Peru, yielding a PeCV DNA prevalence of 15.5 % (9/58). The nested PCR screening for human-associated PeCV DNA was extended to feces from children with diarrhea of unknown etiology from South and Central America [2, 15, 17, 29]. PeCV DNA was found with a detection rate of 5.9 % (3/51) in Nicaraguan and of 3 % (3/100) in Chilean fecal samples. However, none of the diarrhea feces from Brazil (n=170) and Venezuela (n=40) were positive for PeCV DNA. Two PeCV DNA positives (one from Nicaragua and another from Chile) were selected to determine their complete genome sequences by inverse PCR and Sanger sequencing, generating Nicaraguan pecovirus (PeCV-NI, 2928 bases; GenBank KT600066), and Chilean pecovirus (PeCV-CH, 2928 bases; GenBank KT600067). All human-feces-derived genomes were closely related in their Rep gene and more divergent in their Cap genes, with Rep proteins sharing 99 % identity, while the Cap proteins of PeCV-NI and PeCV-CH showed only 76 % identity to that of PeCV-PE. Similar relationships were seen by phylogenetic analysis (Fig. 1B) and genome sequence alignments (Fig. 1C).

Highly divergent small circular DNA genomes

During the initial viral metagenomics analysis, we also found eight sequence reads from two sample pools (four reads per pool) showing similarities to a chimpanzee-feces-associated CRESS-DNA genome (ChiSCV-GM488; GenBank GQ351272). Using iPCR, we determined the complete sequences of these two genomes (Fig. 2). We provisionally called these and related viruses “smacoviruses” (SmaCV2 and SmaCV3) [14]. The genomes of SmaCV2 (2601 bases, GenBank KT600068) and SmaCV3 (2,541 base; GenBank KT600069) also contained two major ORFs encoding Rep and Cap proteins in opposite orientation. The Rep (278 aa) and Cap (369 aa) of SmaCV2 showed the highest identities of 57 % to Rep of chimpanzee-feces-associated ChiSCV-GM488 (GenBank GQ351272) and 50 % to Cap of porcine feces associated SmaCV6 (GenBank KJ577819). The Rep (267 aa) of SmaCV3 showed 95 % identity to captive-chimpanzee-feces-derived smacovirus SF2 (GenBank KP233190),



◀ **Fig. 1** New pecoviruses. **A.** Genome organizations and stem-loop structures. **B.** Phylogenetic trees generated based on Rep and Cap protein sequences from new pecoviruses detected in patients with unexplained diarrhea in Peru, Nicaragua, and Chile and other genetically related CRESS-DNA genomes. The scale indicates amino acid substitutions per position. **C.** Nucleotide sequence similarity among the new pecoviruses. The Rep and Cap locations are shown

while their Cap proteins were 74 % identical. A large stem-loop containing a conserved nonamer (NANT(A/G)TTAC), a small adjoining stem-loop (TAAA), rolling circle replication motifs I-III, and ATP-dependent helicase motifs A-C [19] were detected in SmaCV2 and -3. Phylogenetic analysis showed SmaCV2-3 to be distinct from the other recently described human feces-associated smacoviruses (Fig. 2).

Nested PCR screening did not find additional SmaCV2-positive samples among the 58 Peruvian fecal samples. SmaCV3 DNA could be nPCR amplified from another 10 Peruvian fecal samples, yielding a detection rate of 1.7 % (1/58) and 19 % (11/58) for SmaCV2 and -3, respectively in Peru. None of the fecal samples from children with diarrhea of unknown etiology from South and Central America, including Nicaragua (n=51) [2], Brazil (n=170) [17] and Venezuela (n=40) [29], were SmaCV2 or -3 positive by PCR.

Individual metagenomics analysis of CRESS-DNA containing fecal samples

To investigate whether other viral pathogens were present in the pecovirus- and smacovirus-containing fecal samples from diarrheic patients, five PCR-positive samples from which whole CRESS-DNA genomes were derived were individually analyzed by deep sequencing. As expected, the PeCV sequences were found in the individual feces from Peru (4,704 of 6,971,262 total reads), Nicaragua (537 of 4,704,310 total reads) and Chile (32,603 of 13,458,436 total reads). The following enteric pathogens were also found in the PeCV-positive samples: group A rotavirus (150,301 reads) in the Peruvian sample, enterovirus A (124 reads) in the Nicaraguan sample and enterovirus B (99,112 reads) in the Chilean sample. Deep sequencing also confirmed the presence of SmaCV2 (100 of 448,892 total reads) and SmaCV3 (16 of 770,660 total reads). The

SmaCV2- and SmaCV3-positive feces from Peru both showed the presence of nucleic acids from picobirnavirus (134 and 948 reads). The presence of known enteric pathogens indicates that the diarrhea experienced by the sample donors could be explained by these mammalian viruses rather than the CRESS-DNA viruses.

Discussion

Small circular ssDNA viral genomes were genetically characterized in fecal samples from cases of diarrhea of unknown etiology. We provisionally called these viruses “pecoviruses” and “smacoviruses” [14]. Pecoviruses currently include five distinct genomes from the feces of humans, seals, pigs and dromedaries. The first human-feces-associated PeCV was reported in a patient during an outbreak of acute gastroenteritis in the Netherlands (VS6600022) [27]. Other related genomes were found in feces from asymptomatic mammals [4, 24, 30]. Here, we also report PeCV genomes in human feces from Peru, Nicaragua and Chile, while diarrhea samples from Brazil and Venezuela were PCR negative. All PeCVs shared a similar genome organization, but the conserved nonamer located in a stem-loop differed between human-feces-associated PeCVs (containing a nonamer TTTTATGAG) and animal-feces-derived PeCVs (containing the canonical nonamer TAGTATTAC). The genetic characterization of three related PeCV genomes in human feces from Peru, Nicaragua and Chile plus the previously reported Dutch genome showed highly divergent capsid genes, indicating the presence of diverse PeCVs in human feces. Sequencing of SmaCV2 and -3 also indicated that human-feces-associated SmaCVs are highly diverse, and when compared to other smacoviruses, they exhibited a range of stem-loop structures.

Known or potential enteric viral pathogens were also detected in these diarrheic donors’ feces, which may have played more-direct roles in these patients’ diarrhea. As the cell tropism of these CRESS-DNA genomes remains unknown, so do their roles, if any, in these patients’ symptoms. Detection of these genomes in feces could reflect replication in human gut cells, passive transit of viruses in the food consumed by these individuals, or conceivably, even replication in gut-residing protozoa, fungi, or bacteria.




Fig. 2 New smacoviruses. **A.** Genome organizations and stem-loop structures. **B.** Phylogenetic trees generated based on Rep and Cap protein sequences from smacoviruses detected in Peruvian patients with unexplained diarrhea and other genetically related smacoviruses.

The scale indicates amino acid substitutions per position. **C.** Nucleotide similarity between Peruvian smacoviruses and chimpanzee smacovirus. The Rep and Cap locations are shown

Acknowledgements We acknowledge NHLBI Grant R01 HL105770 to E.L.D and the Blood Systems Research Institute for support. A.C.C. was supported by Sao Paulo Research Foundation (FAPESP), Grant numbers 2014/05211-2 and 2012/03417-7.

Compliance with ethical standards

Funding This study was funded by NHLBI Grant R01 HL105770 to E.L.D and by Sao Paulo Research Foundation (FAPESP) Grant numbers 2014/05211-2 and 2012/03417-7 to A.C.C.

Conflict of interest All authors declare that there is no conflict of interest.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent Pre-existing and anonymized clinical samples were used.

References

- Blinkova O, Victoria J, Li Y, Keele BF, Sanz C, Ndjanga JB, Peeters M, Travis D, Lonsdorf EV, Wilson ML, Pusey AE, Hahn BH, Delwart EL (2010) Novel circular DNA viruses in stool samples of wild-living chimpanzees. *J Gen Virol* 91:74–86
- Bucardo F, Nordgren J, Carlsson B, Paniagua M, Lindgren PE, Espinoza F, Svensson L (2008) Pediatric norovirus diarrhea in Nicaragua. *J Clin Microbiol* 46:2573–2580
- Cheung AK, Ng TF, Lager KM, Bayles DO, Alt DP, Delwart EL, Pogradichny RM, Kehrl ME Jr (2013) A divergent clade of circular single-stranded DNA viruses from pig feces. *Arch Virol* 158:2157–2162
- Cheung AK, Ng TF, Lager KM, Alt DP, Delwart EL, Pogradichny RM (2014) Unique circovirus-like genome detected in pig feces. *Genome Announc* 2(2):e00251–14. doi:10.1128/genomeA.00251-14
- Cheung AK, Ng TF, Lager KM, Alt DP, Delwart E, Pogradichny RM (2015) Identification of several clades of novel single-stranded circular DNA viruses with conserved stem-loop structures in pig feces. *Arch Virol* 160:353–358
- Dayaram A, Goldstien S, Arguello-Astorga GR, Zawar-Reza P, Gomez C, Harding JS, Varsani A (2015) Diverse small circular DNA viruses circulating amongst estuarine molluscs. *Infect Genet Evol J Mol Epidemiol Evol Genet Infect Dis* 31:284–295
- Delwart E, Li L (2012) Rapidly expanding genetic diversity and host range of the Circoviridae viral family and other Rep encoding small circular ssDNA genomes. *Virus Res* 164:114–121
- Espejo PW, Peralta FO, Pacheres HC, del Valle LJ, Tapia AC, Mayra JB, Ruiz J, Mendoza Jdel V (2014) Diarrhoea caused by rotavirus in a regional Peruvian hospital: determination of circulating genotypes. *Trans R Soc Trop Med Hyg* 108:425–430
- Garigliany MM, Borstler J, Jost H, Badusche M, Desmecht D, Schmidt-Chanasit J, Cadar D (2015) Characterization of a novel circovirus-like virus in *Aedes vexans* mosquitoes from Germany: evidence for a new genus within the family Circoviridae. *J Gen Virol* 96:915–920
- Kim HK, Park SJ, Nguyen VG, Song DS, Moon HJ, Kang BK, Park BK (2012) Identification of a novel single-stranded, circular DNA virus from bovine stool. *J Gen Virol* 93:635–639
- Krabberger S, Arguello-Astorga GR, Greenfield LG, Galilee C, Law D, Martin DP, Varsani A (2015) Characterisation of a diverse range of circular replication-associated protein encoding DNA viruses recovered from a sewage treatment oxidation pond. *Infect Genet Evol J Mol Epidemiol Evol Genet Infect Dis* 31:73–86
- Labonte JM, Suttle CA (2013) Previously unknown and highly divergent ssDNA viruses populate the oceans. *ISME J* 7:2169–2177
- Ng TF, Marine R, Wang C, Simmonds P, Kapusinszky B, Bodhidatta L, Oderinde BS, Wommack KE, Delwart E (2012) High variety of known and new RNA and DNA viruses of diverse origins in untreated sewage. *J Virol* 86:12161–12175
- Ng TFF, Zhang W, Sachsenroder J, Kondov NO, Antonio Charlyes da Costa, Vega E, Holtz LR, Wu G, Wang D, Stine CO, Antonio M, Mulvaney US, Muench MO, Deng X, Ambert-Balay K, Pothier P, Vinje J, Delwart E (2015) A diverse group of small circular ssDNA viral genomes in human and non-human primate stools. *Virus Evo* 1 (1): 1–3. doi:10.1093/ve/vev017
- Phan TG, Li L, O’Ryan MG, Cortes H, Mamani N, Bonkougou IJ, Wang C, Leutenegger CM, Delwart E (2012) A third gyrovirus species in human faeces. *J Gen Virol* 93:1356–1361
- Phan TG, Vo NP, Boros A, Pankovics P, Reuter G, Li OT, Wang C, Deng X, Poon LL, Delwart E (2013) The viruses of wild pigeon droppings. *PLoS ONE* 8:e72787
- Phan TG, Mori D, Deng X, Rajindrajith S, Ranawaka U, Fan Ng TF, Bucardo-Rivera F, Orlandi P, Ahmed K, Delwart E (2015) Small circular single stranded DNA viral genomes in unexplained cases of human encephalitis, diarrhea, and in untreated sewage. *Virology* 482:98–104
- Reuter G, Boros A, Delwart E, Pankovics P (2014) Novel circular single-stranded DNA virus from turkey faeces. *Arch Virol* 159:2161–2164
- Rosario K, Duffy S, Breitbart M (2012) A field guide to eukaryotic circular single-stranded DNA viruses: insights gained from metagenomics. *Arch Virol* 157:1851–1871
- Rosario K, Schenck RO, Harbeitner RC, Lawler SN, Breitbart M (2015) Novel circular single-stranded DNA viruses identified in marine invertebrates reveal high sequence diversity and consistent predicted intrinsic disorder patterns within putative structural proteins. *Front Microbiol* 6:696
- Sachsenroder J, Twardziok S, Hammerl JA, Janczyk P, Wrede P, Hertwig S, Johne R (2012) Simultaneous identification of DNA and RNA viruses present in pig faeces using process-controlled deep sequencing. *PLoS ONE* 7:e34631
- Sachsenroder J, Braun A, Machnowska P, Ng TF, Deng X, Guenther S, Bernstein S, Ulrich RG, Delwart E, Johne R (2014) Metagenomic identification of novel enteric viruses in urban wild rats and genome characterization of a group A rotavirus. *J Gen Virol* 95:2734–2747
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
- Sikorski A, Dayaram A, Varsani A (2013) Identification of a novel circular DNA virus in New Zealand Fur Seal (*Arctocepalhus forsteri*) fecal matter. *Genome Announc* 1(4):e00558–13. doi:10.1128/genomeA.00558-13
- Sikorski A, Massaro M, Kraberger S, Young LM, Smalley D, Martin DP, Varsani A (2013) Novel myco-like DNA viruses discovered in the faecal matter of various animals. *Virus Res* 177:209–216
- Smits SL, Raj VS, Oduber MD, Schapendonk CM, Bodewes R, Provacia L, Stittelaar KJ, Osterhaus AD, Haagmans BL (2013) Metagenomic analysis of the ferret fecal viral flora. *PLoS ONE* 8:e71595
- Smits SL, Schapendonk CM, van Beek J, Vennema H, Schurch AC, Schipper D, Bodewes R, Haagmans BL, Osterhaus AD, Koopmans MP (2014) New viruses in idiopathic human diarrhea cases, the Netherlands. *Emerg Infect Dis* 20:1218–1222
- Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S (2013) MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30:2725–2729

29. [Vizzi E, Pineros O, Gonzalez GG, Zambrano JL, Ludert JE, Liprandi F \(2011\) Genotyping of human rotaviruses circulating among children with diarrhea in Valencia, Venezuela. J Med Virol 83:2225–2232](#)
30. [Woo PC, Lau SK, Teng JL, Tsang AK, Joseph M, Wong EY, Tang Y, Sivakumar S, Bai R, Wernery R, Wernery U, Yuen KY \(2014\) Metagenomic analysis of viromes of dromedary camel fecal samples reveals large number and high diversity of circoviruses and picobirnaviruses. Virology 471–473:117–125](#)
31. [Yu X, Li B, Fu Y, Jiang D, Ghabrial SA, Li G, Peng Y, Xie J, Cheng J, Huang J, Yi X \(2010\) A geminivirus-related DNA mycovirus that confers hypovirulence to a plant pathogenic fungus. Proc Natl Acad Sci USA 107:8387–8392](#)
32. [Zawar-Reza P, Arguello-Astorga GR, Kraberger S, Julian L, Stainton D, Broady PA, Varsani A \(2014\) Diverse small circular single-stranded DNA viruses identified in a freshwater pond on the McMurdo Ice Shelf \(Antarctica\). Infect Genet Evol J Mol Epidemiol Evol Genet Infect Dis 26:132–138](#)
33. [Zuker M \(2003\) Mfold web server for nucleic acid folding and hybridization prediction. Nucleic Acids Res 31:3406–3415](#)